


## Multi-configurational self-consistent field

*Emmanuel Fromager*



Institut de Chimie de Strasbourg - Laboratoire de Chimie Quantique -  
Université de Strasbourg /CNRS

ISTPC 2017 summer school, Aussois, France.



## Notations

- Molecular orbitals:  $\phi_p(\mathbf{r}) = \sum_{\mu} C_{\mu p} \chi_{\mu}(\mathbf{r})$   $\langle \phi_p | \phi_q \rangle = \delta_{pq}$
- Non-orthogonal set of atomic orbitals (Gaussian functions):  $\langle \chi_{\mu} | \chi_{\nu} \rangle = S_{\mu\nu}$
- Hamiltonian in second quantization:

$$\hat{H} = \sum_{p,q} h_{pq} \hat{E}_{pq} + \frac{1}{2} \sum_{p,q,r,s} \langle pr | qs \rangle \left( \hat{E}_{pq} \hat{E}_{rs} - \delta_{qr} \hat{E}_{ps} \right)$$

where 
$$h_{pq} = \int d\mathbf{r} \phi_p(\mathbf{r}) \left[ -\frac{1}{2} \nabla_{\mathbf{r}}^2 + v_{\text{ne}}(\mathbf{r}) \right] \phi_q(\mathbf{r})$$

and 
$$\langle pr | qs \rangle = \iint d\mathbf{r}_1 d\mathbf{r}_2 \phi_p(\mathbf{r}_1) \phi_r(\mathbf{r}_2) \frac{1}{|\mathbf{r}_1 - \mathbf{r}_2|} \phi_q(\mathbf{r}_1) \phi_s(\mathbf{r}_2) = (pq | rs)$$

## Variational and non-variational approximations

- The **exact** electronic ground state  $\Psi_0$  and its energy  $E_0$  can be obtained two ways:

$$E_0 = \min_{\Psi} \frac{\langle \Psi | \hat{H} | \Psi \rangle}{\langle \Psi | \Psi \rangle} = \frac{\langle \Psi_0 | \hat{H} | \Psi_0 \rangle}{\langle \Psi_0 | \Psi_0 \rangle} \quad \hat{H} | \Psi_0 \rangle = E_0 | \Psi_0 \rangle$$

- Approximate parametrized** ground-state wave function:  $\Psi(\boldsymbol{\lambda}_0)$

where  $\boldsymbol{\lambda}_0$  denotes the complete set of **optimized** parameters.

### Variational calculation

$$\left. \frac{\partial}{\partial \boldsymbol{\lambda}} \frac{\langle \Psi(\boldsymbol{\lambda}) | \hat{H} | \Psi(\boldsymbol{\lambda}) \rangle}{\langle \Psi(\boldsymbol{\lambda}) | \Psi(\boldsymbol{\lambda}) \rangle} \right|_{\boldsymbol{\lambda}=\boldsymbol{\lambda}_0} = 0$$

↓

Hartree-Fock (HF)

Configuration Interaction (CI)

Multi-Configurational Self-Consistent Field (MCSCF)

### Non-variational calculation

$$\hat{H} | \Psi(\boldsymbol{\lambda}) \rangle - E(\boldsymbol{\lambda}) | \Psi(\boldsymbol{\lambda}) \rangle = 0 \quad \text{for } \boldsymbol{\lambda} = \boldsymbol{\lambda}_0$$

↓

Many-Body Perturbation Theory (MBPT)

Coupled Cluster (CC)

## Spin-orbital rotation

- Let  $\{\varphi_P\}_P$  denote an orthonormal basis of spin-orbitals and  $\{\tilde{\varphi}_P\}_P$  another orthonormal basis obtained by **unitary transformation**:

$$|\tilde{\varphi}_P\rangle = \sum_Q U_{QP} |\varphi_Q\rangle$$

- U can be written as  $\boxed{U = e^{-\kappa} \text{ with } \kappa^\dagger = -\kappa}$   $\leftarrow U^\dagger = (e^{-\kappa})^\dagger = e^{-\kappa^\dagger} = e^\kappa = U^{-1}$
- $\kappa_{PQ}$  can be used rather than  $U_{PQ}$  for parametrizing the spin-orbital rotation

**EX1:** Using EX2, show that in **second quantization** the unitary transformation can be simply written as

$$\hat{a}_{\tilde{P}}^\dagger = \sum_Q (e^{-\kappa})_{QP} \hat{a}_Q^\dagger = \boxed{e^{-\hat{\kappa}} \hat{a}_P^\dagger e^{\hat{\kappa}} = \hat{a}_{\tilde{P}}^\dagger} \quad \text{where} \quad \hat{\kappa} = \sum_{PQ} \kappa_{PQ} \hat{a}_P^\dagger \hat{a}_Q$$

## Spin-orbital rotation

- Note that the rotation operator  $\hat{\kappa}$  is anti-Hermitian:

$$\hat{\kappa}^\dagger = \sum_{PQ} \kappa_{PQ}^* \hat{a}_Q^\dagger \hat{a}_P = \sum_{PQ} \kappa_{QP}^\dagger \hat{a}_Q^\dagger \hat{a}_P = - \sum_{PQ} \kappa_{QP} \hat{a}_Q^\dagger \hat{a}_P = -\hat{\kappa}$$

- Unitary transformation for a  $N$ -electron Slater determinant:

$$\begin{aligned} |\tilde{P}_1 \tilde{P}_2 \dots \tilde{P}_N\rangle &= \hat{a}_{\tilde{P}_1}^\dagger \hat{a}_{\tilde{P}_2}^\dagger \dots \hat{a}_{\tilde{P}_N}^\dagger |\text{vac}\rangle = e^{-\hat{\kappa}} \hat{a}_{P_1}^\dagger e^{\hat{\kappa}} e^{-\hat{\kappa}} \hat{a}_{P_2}^\dagger e^{\hat{\kappa}} \dots e^{-\hat{\kappa}} \hat{a}_{P_N}^\dagger e^{\hat{\kappa}} |\text{vac}\rangle \\ &= e^{-\hat{\kappa}} \hat{a}_{P_1}^\dagger \hat{a}_{P_2}^\dagger \dots \hat{a}_{P_N}^\dagger \underbrace{e^{\hat{\kappa}} |\text{vac}\rangle}_{|\text{vac}\rangle} \end{aligned}$$

$$|\tilde{P}_1 \tilde{P}_2 \dots \tilde{P}_N\rangle = e^{-\hat{\kappa}} |P_1 P_2 \dots P_N\rangle$$

## Spin-restricted orbital rotation

- In a **restricted formalism** the same set of orbitals is used for  $\alpha$  and  $\beta$  spins:

$$\hat{\kappa} = \sum_{PQ} \kappa_{PQ} \hat{a}_P^\dagger \hat{a}_Q = \sum_{pq} \sum_{\sigma\sigma'} \underbrace{\kappa_{p,\sigma q,\sigma'}}_{\kappa_{pq} \delta_{\sigma\sigma'}} \hat{a}_{p,\sigma}^\dagger \hat{a}_{q,\sigma'} = \sum_{pq} \kappa_{pq} \hat{E}_{pq}$$

- Since  $\kappa_{pq} = -\kappa_{qp}$  (real algebra)

$$\hat{\kappa} = \sum_{p>q} \kappa_{pq} \hat{E}_{pq} - \sum_{p<q} \kappa_{qp} \hat{E}_{pq}$$

$$\hat{\kappa} = \sum_{p>q} \kappa_{pq} (\hat{E}_{pq} - \hat{E}_{qp})$$

### Hartree-Fock approximation

- For simplicity we consider here the particular case of a non-degenerate **singlet closed-shell** ground state
- The HF method consists then in **approximating** the exact **wave function**  $\Psi_0$  by a single **Slater determinant**  $\Phi_0$ . The orbital space is thus divided in two:

**doubly occupied molecular orbitals**  $\phi_i, \phi_j, \dots$

**unoccupied molecular orbitals**  $\phi_a, \phi_b, \dots$

$$|\Phi_0\rangle = \prod_i^{\text{occ.}} \prod_{\sigma=\alpha,\beta} \hat{a}_{i,\sigma}^\dagger |\text{vac}\rangle$$

- The initial set of molecular orbitals is usually not optimized  $\rightarrow$  the **optimized** HF molecular orbitals will be obtained by means of **unitary transformations** (orbital rotation)



## Hartree-Fock approximation

- Exponential parametrization:  $|\Phi(\boldsymbol{\kappa})\rangle = e^{-\hat{\kappa}} |\Phi_0\rangle$  with  $\hat{\kappa} = \sum_{p>q} \kappa_{pq} (\hat{E}_{pq} - \hat{E}_{qp})$

$$\boldsymbol{\kappa} = \begin{bmatrix} \vdots \\ \kappa_{pq} \\ \vdots \end{bmatrix}_{p>q}$$

denotes the column vector containing **all the parameters to be optimized**

- occupied-occupied and unoccupied-unoccupied rotations:

$$\hat{\kappa} = \underbrace{\sum_{i>j} \kappa_{ij} (\hat{E}_{ij} - \hat{E}_{ji})}_{\hat{\kappa}^{\text{occ.}}} + \sum_{i,a} \kappa_{ai} (\hat{E}_{ai} - \hat{E}_{ia}) + \underbrace{\sum_{a>b} \kappa_{ab} (\hat{E}_{ab} - \hat{E}_{ba})}_{\hat{\kappa}^{\text{unocc.}}}$$

$$\hat{\kappa}^{\text{occ.}} |\Phi_0\rangle = \hat{\kappa}^{\text{unocc.}} |\Phi_0\rangle = 0 \rightarrow \text{only **occupied-unoccupied rotations** will be optimized} \rightarrow \boldsymbol{\kappa} = \begin{bmatrix} \vdots \\ \kappa_{ai} \\ \vdots \end{bmatrix}$$

## Hartree-Fock approximation

- Hartree-Fock energy expression:

$$E(\boldsymbol{\kappa}) = \frac{\langle \Phi(\boldsymbol{\kappa}) | \hat{H} | \Phi(\boldsymbol{\kappa}) \rangle}{\langle \Phi(\boldsymbol{\kappa}) | \Phi(\boldsymbol{\kappa}) \rangle} = \frac{\langle \Phi_0 | e^{-\hat{\kappa}^\dagger} \hat{H} e^{-\hat{\kappa}} | \Phi_0 \rangle}{\langle \Phi_0 | e^{-\hat{\kappa}^\dagger} e^{-\hat{\kappa}} | \Phi_0 \rangle} = \boxed{\langle \Phi_0 | e^{\hat{\kappa}} \hat{H} e^{-\hat{\kappa}} | \Phi_0 \rangle = E(\boldsymbol{\kappa})}$$

- Variational optimization of  $\boldsymbol{\kappa}$ :  $E_{\boldsymbol{\kappa}_+}^{[1]} = \left. \frac{\partial E(\boldsymbol{\kappa})}{\partial \boldsymbol{\kappa}} \right|_{\boldsymbol{\kappa}_+} = 0$

- Iterative procedure (Newton method):

$$E(\boldsymbol{\kappa}) \approx E(0) + \boldsymbol{\kappa}^T E_0^{[1]} + \frac{1}{2} \boldsymbol{\kappa}^T E_0^{[2]} \boldsymbol{\kappa} \rightarrow E_{\boldsymbol{\kappa}_+}^{[1]} \approx E_0^{[1]} + E_0^{[2]} \boldsymbol{\kappa}_+ = 0 \rightarrow E_0^{[2]} \underbrace{\boldsymbol{\kappa}_+}_{\text{Newton step}} = -E_0^{[1]}$$

- Update the HF determinant:  $\Phi_0 \leftarrow \Phi(\boldsymbol{\kappa}_+)$

Newton step

- HF calculation converged when  $\boxed{E_0^{[1]} = 0}$

## Analytical gradient and Hessian in Hartree-Fock

**EX2:** By using the relation  $e^{-\lambda\hat{A}} \hat{B} e^{\lambda\hat{A}} = \hat{B} + \int_0^\lambda d\xi \frac{d}{d\xi} \left( e^{-\xi\hat{A}} \hat{B} e^{\xi\hat{A}} \right)$ , which holds for any  $\lambda$  and any  $\hat{B}$  operator, prove the *Baker-Campbell-Hausdorff* (BCH) expansion:

$$e^{-\hat{A}} \hat{B} e^{\hat{A}} = \hat{B} + \sum_{n=1}^{+\infty} \frac{1}{n!} [[\hat{B}, \hat{A}]_n$$

$$[[\hat{B}, \hat{A}]_{n+1} = [[\hat{B}, \hat{A}]_n, \hat{A}], \quad [[\hat{B}, \hat{A}]_1 = [\hat{B}, \hat{A}]$$


$$= \hat{B} + [\hat{B}, \hat{A}] + \frac{1}{2} [[\hat{B}, \hat{A}], \hat{A}] + \dots$$

- **Analytical formulas** for the **gradient** and the **hessian**:


$$E(\boldsymbol{\kappa}) = E(0) + \underbrace{\langle \Phi_0 | [\hat{\boldsymbol{\kappa}}, \hat{H}] | \Phi_0 \rangle}_{\text{gradient}} + \frac{1}{2} \langle \Phi_0 | [\hat{\boldsymbol{\kappa}}, [\hat{\boldsymbol{\kappa}}, \hat{H}]] | \Phi_0 \rangle + \dots$$

$$\sum_{ai} \kappa_{ai} \langle \Phi_0 | [\hat{E}_{ai} - \hat{E}_{ia}, \hat{H}] | \Phi_0 \rangle \quad \rightarrow \quad E_{0,ai}^{[1]} = \langle \Phi_0 | [\hat{E}_{ai} - \hat{E}_{ia}, \hat{H}] | \Phi_0 \rangle$$


$$= -2 \langle \Phi_0 | \hat{H} \hat{E}_{ai} | \Phi_0 \rangle = 0 \quad (\text{Brillouin theorem})$$



Short-range dynamical correlation



Static correlation



## Static correlation

- H<sub>2</sub> in the equilibrium geometry:

$$\boxed{|\Psi_0\rangle = C_0|1\sigma_g^\alpha 1\sigma_g^\beta\rangle + \dots} \quad \text{where } |C_0|^2 = 98\% \quad \text{no static correlation}$$


- In the dissociation limit: H<sub>A</sub>...H<sub>B</sub> and **NOT** H<sub>A</sub><sup>-</sup>...H<sub>B</sub><sup>+</sup> or H<sub>A</sub><sup>+</sup>...H<sub>B</sub><sup>-</sup>

$$\phi_{1\sigma_g}(\mathbf{r}) = \frac{1}{\sqrt{2}} \left( \phi_{1s_A}(\mathbf{r}) + \phi_{1s_B}(\mathbf{r}) \right) \quad \text{and} \quad \phi_{1\sigma_u}(\mathbf{r}) = \frac{1}{\sqrt{2}} \left( \phi_{1s_A}(\mathbf{r}) - \phi_{1s_B}(\mathbf{r}) \right)$$

$$|1\sigma_g^\alpha 1\sigma_g^\beta\rangle = \frac{1}{2} \left( |1s_A^\alpha 1s_B^\beta\rangle + |1s_B^\alpha 1s_A^\beta\rangle + |1s_A^\alpha 1s_A^\beta\rangle + |1s_B^\alpha 1s_B^\beta\rangle \right)$$

$$-|1\sigma_u^\alpha 1\sigma_u^\beta\rangle = \frac{1}{2} \left( |1s_A^\alpha 1s_B^\beta\rangle + |1s_B^\alpha 1s_A^\beta\rangle - |1s_A^\alpha 1s_A^\beta\rangle - |1s_B^\alpha 1s_B^\beta\rangle \right)$$

$$\boxed{|\Psi_0\rangle = \frac{1}{\sqrt{2}} \left( |1\sigma_g^\alpha 1\sigma_g^\beta\rangle - |1\sigma_u^\alpha 1\sigma_u^\beta\rangle \right)} \quad \text{strong static correlation}$$





### Multi-Configurational Self-Consistent Field model (MCSCF)

- The MCSCF model consists in performing a CI calculation with a **reoptimization of the orbitals**

$$|\Psi(\boldsymbol{\kappa}, \mathbf{C})\rangle = e^{-\hat{\kappa}} \left( \sum_{\xi} C_{\xi} |\text{det}_{\xi}\rangle \right)$$

- The MCSCF model is a **multiconfigurational extension of HF** which aims at describing **static correlation**: a limited number of determinants should be sufficient.
- Short-range dynamical correlation is treated afterwards (post-MCSCF models)
- Choice of the determinants: **active space**

H. . .H                      2 electrons in 2 orbitals ( $1\sigma_g, 1\sigma_u$ )                       $\longrightarrow$  2/2

Be                              2 electrons in 4 orbitals ( $2s, 2p_x, 2p_y, 2p_z$ )                       $\longrightarrow$  2/4

## Multi-Configurational Self-Consistent Field model (MCSCF)

- Complete Active Space (**CAS**) for Be:  $|1s^2 2s^2\rangle, |1s^2 2p_x^2\rangle, |1s^2 2p_y^2\rangle, |1s^2 2p_z^2\rangle,$

if all the determinants are included in the MCSCF calculation  $\longrightarrow$  CASSCF

if a Restricted Active Space (**RAS**) is used  $\longrightarrow$  RASSCF

- The orbital space is now divided in three:

doubly occupied molecular orbitals (inactive)	$\phi_i, \phi_j, \dots$	$1s$
<b>active</b> molecular orbitals	$\phi_u, \phi_v, \dots$	$2s, 2p_x, 2p_y, 2p_z$
unoccupied molecular orbitals	$\phi_a, \phi_b, \dots$	$3s, 3p, 3d, \dots$

# Multi-configurational self-consistent field


$a, b, \dots$




$u, v, \dots$



$i, j, \dots$



$H_2$  (FCI, aug-cc-pVQZ)

- Iterative optimization of the orbital rotation vector  $\kappa$  and the CI coefficients  $C_i$ :

$$|\Psi^{(0)}\rangle = \sum_i C_i^{(0)} |i\rangle$$

← **normalized** starting wave function

$$|\Psi(\lambda)\rangle = e^{-\hat{\kappa}} \frac{|\Psi^{(0)}\rangle + \hat{Q}|\delta\rangle}{\sqrt{1 + \langle \delta | \hat{Q} | \delta \rangle}}$$

← convenient parametrization  $\lambda =$

$$\begin{bmatrix} \vdots \\ \kappa_{pq} \\ \vdots \\ \delta_i \\ \vdots \end{bmatrix} \quad p > q$$

$$\hat{Q} = 1 - |\Psi^{(0)}\rangle\langle\Psi^{(0)}|, \quad |\delta\rangle = \sum_i \delta_i |i\rangle, \quad \langle\Psi^{(0)}|\hat{Q}|\delta\rangle = 0, \quad \langle\Psi(\lambda)|\Psi(\lambda)\rangle = 1$$

- MCSCF energy expression:  $E(\lambda) = \langle\Psi(\lambda)|\hat{H}|\Psi(\lambda)\rangle$

- Variational optimization:

$$E_{\lambda_+}^{[1]} = \begin{bmatrix} E_{\lambda_+}^{o[1]} \\ E_{\lambda_+}^{c[1]} \end{bmatrix} = 0$$

where  $E_{\lambda_+}^{o[1]} = \left. \frac{\partial E(\lambda)}{\partial \kappa} \right|_{\lambda_+}$

and  $E_{\lambda_+}^{c[1]} = \left. \frac{\partial E(\lambda)}{\partial \delta} \right|_{\lambda_+}$

- Newton method:

$$E(\boldsymbol{\lambda}) \approx E(0) + \boldsymbol{\lambda}^T E_0^{[1]} + \frac{1}{2} \boldsymbol{\lambda}^T E_0^{[2]} \boldsymbol{\lambda} \quad \rightarrow \quad E_{\boldsymbol{\lambda}_+}^{[1]} \approx E_0^{[1]} + E_0^{[2]} \boldsymbol{\lambda}_+ = 0 \quad \rightarrow \quad E_0^{[2]} \underbrace{\boldsymbol{\lambda}_+}_{\text{Newton step}} = -E_0^{[1]}$$


**Newton step**

- Convergence reached when  $E_0^{[1]} = 0$

**EX7:** Show that  $E_{0,pq}^{o[1]} = \langle \Psi^{(0)} | [\hat{E}_{pq} - \hat{E}_{qp}, \hat{H}] | \Psi^{(0)} \rangle$  and  $E_0^{c[1]} = 2 \left( \mathbf{H}^{\text{CAS}} - E(0) \right) \mathbf{C}^{(0)}$

where  $\mathbf{H}_{ij}^{\text{CAS}} = \langle i | \hat{H} | j \rangle$  and  $\mathbf{C}^{(0)} = \begin{bmatrix} \vdots \\ C_i^{(0)} \\ \vdots \end{bmatrix}$

Note:  $E_0^{o[1]} = 0$  is known as **generalized Brillouin theorem**.



## Multi-state MCSCF approach

- **State-averaged** MCSCF model: **simultaneous optimization** of the ground and the lowest  $\mathcal{N} - 1$  excited states at the MCSCF level.
- Iterative procedure:  $\mathcal{N}$  initial orthonormal states are built from the same set of orbitals.

$$|\Psi_I^{(0)}\rangle = \sum_i C_{I,i}^{(0)} |i\rangle, \quad I = 1, \dots, \mathcal{N}$$

- **Double-exponential** parametrization:

$$\boxed{|\Psi_I(\boldsymbol{\kappa}, \mathbf{S})\rangle = e^{-\hat{\kappa}} e^{-\hat{S}} |\Psi_I^{(0)}\rangle} \quad \text{where} \quad \hat{S} = \sum_{J=1}^{\mathcal{N}} \sum_{K>J} S_{KJ} \left( |\Psi_K^{(0)}\rangle \langle \Psi_J^{(0)}| - |\Psi_J^{(0)}\rangle \langle \Psi_K^{(0)}| \right)$$

$$\text{and} \quad \sum_i |i\rangle \langle i| = \sum_K |\Psi_K^{(0)}\rangle \langle \Psi_K^{(0)}|$$



## Multi-state MCSCF approach

- Gross–Oliveira–Kohn (GOK) **variational principle** for an ensemble of **ground and excited states**:

For any set  $\{\Psi_I\}_{I=1,\mathcal{N}}$  of  $\mathcal{N}$  orthonormal states, the following inequality holds,

$$\sum_{I=1}^{\mathcal{N}} w_I \langle \Psi_I | \hat{H} | \Psi_I \rangle \geq \sum_{I=1}^{\mathcal{N}} w_I E_I$$

where  $E_1 \leq E_2 \leq \dots \leq E_{\mathcal{N}}$  are the  $\mathcal{N}$  lowest **exact** eigenvalues of  $\hat{H}$ , and the weights are ordered as follows,

$$w_1 \geq w_2 \geq \dots \geq w_{\mathcal{N}} > 0.$$

**EXERCISE:** Prove the theorem in the particular case of two states by using **Theophilou's variational principle**:  $\langle \Psi_1 | \hat{H} | \Psi_1 \rangle + \langle \Psi_2 | \hat{H} | \Psi_2 \rangle \geq E_1 + E_2$ . **Hint:** Show that

$$w_1 \langle \Psi_1 | \hat{H} | \Psi_1 \rangle + w_2 \langle \Psi_2 | \hat{H} | \Psi_2 \rangle = w_2 \left[ \langle \Psi_1 | \hat{H} | \Psi_1 \rangle + \langle \Psi_2 | \hat{H} | \Psi_2 \rangle \right] + (w_1 - w_2) \langle \Psi_1 | \hat{H} | \Psi_1 \rangle$$

**EXERCISE:** Proof of Theophilou's variational principle for two states

(1) Let  $\Delta = \langle \Psi_1 | \hat{H} | \Psi_1 \rangle + \langle \Psi_2 | \hat{H} | \Psi_2 \rangle - E_1 - E_2$ . We consider the complete basis of the exact eigenvectors  $\{ \tilde{\Psi}_I \}_{I=1,2,\dots}$  of  $\hat{H}$  with eigenvalues  $\{ E_I \}_{I=1,2,\dots}$

Both trial wavefunctions can be expanded in that basis as follows,

$$|\Psi_K\rangle = \sum_I C_{KI} |\tilde{\Psi}_I\rangle, \quad K = 1, 2.$$

Show that  $\Delta = \sum_{I=1}^2 (p_I - 1)E_I + \sum_{I>2} p_I E_I$  where  $p_I = C_{1I}^2 + C_{2I}^2$ .

(2) Show that  $\Delta = \sum_{I=1}^2 (1 - p_I)(E_2 - E_I) + \sum_{I>2} p_I (E_I - E_2)$ . **Hint:** prove first that  $\sum_I p_I = 2$ .

(3) Let us now decompose the two first eigenvectors ( $I = 1, 2$ ) in the basis of the trial wavefunctions and the orthogonal complement:  $|\tilde{\Psi}_I\rangle = C_{1I}|\Psi_1\rangle + C_{2I}|\Psi_2\rangle + \hat{Q}_{12}|\tilde{\Psi}_I\rangle$  where

$\hat{Q}_{12} = 1 - \sum_{K=1}^2 |\Psi_K\rangle\langle\Psi_K|$ . Explain why  $p_I \leq 1$  when  $I = 1, 2$  and conclude.

### Multi-state MCSCF approach

- State-averaged energy: 
$$E(\boldsymbol{\kappa}, \mathbf{S}) = \sum_{I=1}^{\mathcal{N}} w_I \langle \Psi_I(\boldsymbol{\kappa}, \mathbf{S}) | \hat{H} | \Psi_I(\boldsymbol{\kappa}, \mathbf{S}) \rangle$$

where  $w_I$  are **arbitrary weights**. In the so-called "equal weight" state-averaged MCSCF calculation  $w_I = \frac{1}{\mathcal{N}}$ .

- Variational optimization: 
$$\frac{\partial E(\boldsymbol{\kappa}, \mathbf{S})}{\partial \boldsymbol{\kappa}} = \frac{\partial E(\boldsymbol{\kappa}, \mathbf{S})}{\partial \mathbf{S}} = 0$$
- Note that, in contrast to the exact theory, converged **individual energies** (and therefore excitation energies) **may vary with the weights**. This is due to the **orbital optimization**.
- **Short-range dynamical correlation** is usually recovered with multi-reference perturbation theory (**multi-state** CASPT2 or NEVPT2 for example) or multi-reference CI.